

Math 319, Fall 2013, Term Test II  
Techniques in Ordinary Differential Equations

Date: Friday, November 15

**Lecture Section: 002**

Name (printed): \_\_\_\_\_

UW Student ID Number: \_\_\_\_\_

Discussion Section: (circle)

<b>Liu Liu:</b>	<b>301</b>	<b>302</b>	<b>303</b>	<b>304</b>
<b>Huanyu Wen:</b>	<b>305</b>	<b>306</b>	<b>323</b>	<b>324</b>
<b>Dongfei Pei:</b>	<b>325</b>	<b>326</b>	<b>329</b>	
<b>Kai Hsu:</b>	<b>327</b>	<b>328</b>		

**Instructions**

1. Fill out this cover page **completely** and make sure to circle your discussion section.
2. Answer questions in the space provided, using the last page for overflow.
3. Show all the work required to obtain your answers.
4. No calculators are permitted.

FOR EXAMINERS' USE ONLY	
Page	Mark
2	/6
3	/7
4	/6
5	/6
Total	/25

1. Short Answer:

- [1] (a) Suppose that  $y_1(x)$  and  $y_2(x)$  are two solutions of the homogeneous second-order differential equation

$$y''(x) + p(x)y'(x) + q(x)y(x) = 0$$

satisfying  $W(y_1, y_2)(x) \neq 0$ . State the general solution form of the differential equation.

- [2] (b) Determine the values  $a$  for which the following differential equations exhibit *oscillatory* behavior:

(i)  $y''(x) + ay'(x) + y(x) = 0$

(ii)  $ay''(x) - 2y'(x) + 2y(x) = 0$

- [3] 2. True/False:

(a) The mechanical mechanism  $x''(t) + cx'(t) + 4x(t) = 0$  is *critically damped* at the value  $c = 4$ . [True / False]

(b) The amplitude (i.e. maximal value) of  $y(x) = 3 \cos(x) - 4 \sin(x)$  is 5. [True / False]

(c) Every point  $x_0 \in \mathbb{R}$  is an ordinary point of  $(x^2 - 1)y''(x) + xy'(x) - (x + 1)y(x) = 0$ . [True / False]

3. Second-Order Differential Equations

- [3] (a) Consider the following differential equation:

$$y''(x) + 6y'(x) + 5y(x) = g(x).$$

Determine the complementary solution  $y_c(x)$  then set-up, but *do not attempt to evaluate*, the trial form  $y_p(x)$  for the given choices of  $g(x)$  below.

(i)  $g(x) = e^{-5x} \sin(x)$

(ii)  $g(x) = xe^{-x}$

- [4] (b) Given that
- $y_1(x) = 1 + x$
- and
- $y_2(x) = e^x$
- are fundamental solutions of
- $xy''(x) - (1 + x)y'(x) + y(x) = 0$
- , determine the particular solution
- $y_p(x)$
- of

$$xy''(x) - (1 + x)y'(x) + y(x) = 2x^2e^x.$$

[*Hint*: Remember the standard form when applying variation of parameters!]

4. Power Series Solutions

- [2] (a) Determine the radius of convergence of the power series:

$$\sum_{n=0}^{\infty} \frac{(-1)^{n+1}(n+1)}{n!} x^n$$

- [2] (b) Write the following expression as a single summation whose common term is  $x^n$ :

$$x \sum_{n=1}^{\infty} n(n+1)a_n x^{n-2} + \sum_{n=2}^{\infty} (n-1)a_n x^{n-2}$$

- [2] (c) Suppose that a particular differential equation has a power series solution centered at  $x_0 = 0$  generated by the recursion relation

$$a_{n+2} = \frac{a_{n+1} + a_n}{(n+2)(n+1)}, \quad n \geq 0.$$

Determine up to the  $x^3$  term of the solution  $y(x) = \sum_{n=0}^{\infty} a_n x^n$ .

5. Laplace Transforms:

- [2] (a) Evaluate the following inverse Laplace transform:

$$\mathcal{L}^{-1} \left\{ \frac{s+2}{(s-1)^2+9} \right\}$$

- [4] (b) Use the Laplace transform method to solve the following initial value problem:

$$2y''(x) - 3y'(x) + y(x) = 0, \quad y(0) = 0, \quad y'(0) = 1/2.$$

THIS PAGE IS FOR ROUGH WORK